



Rhizolizer™

SOIL AMENDMENT

GUARANTEED ANALYSIS

Soil Amending Ingredients:

Microbial Inoculant.....	10.0%
<i>Trichoderma harzianum</i>	1x10 ⁸ CFU/mL
<i>Bacillus amyloliquefaciens</i>	1x10 ⁹ CFU/mL
Total Other Ingredients.....	90.0%

F002455

Not recommended for use as a fertilizer substitute

Emergency Number:
Contact Chemtrec
1(800) 424-9300



For General Information:
Contact Locus Agricultural Solutions
1(888) 331-5008

**KEEP OUT OF REACH OF CHILDREN • SHAKE OR STIR WELL BEFORE USE
FOR PROFESSIONAL, COMMERCIAL, OR INDUSTRIAL USE ONLY**

Density: 8.34 lbs./gallon at 68°F
NET CONTENTS:

- 15 fl. oz. (444 mL)
- 30 fl. oz. (887mL)
- Citrus
- Veg/Fruit
- Sod

LOT #:

Expiration Date:

Manufactured by: **Locus Agricultural Solutions**
30500 Aurora Road, Suite 180
Solon, Ohio 44139

Phone: 1(888) 331-5008 • Copyright © 2018 All rights reserved

PRODUCT INFORMATION/PURPOSE

Rhizolizer™ augments natural microbial activity in the rhizosphere of agricultural soils. It supports nutrient availability and crop nutrient uptake from the soil.

For professional, commercial, or industrial use only.

DIRECTIONS FOR USE

Rhizolizer™ may be applied to soil through drip irrigation, micro-jets, drench, or boom sprayers. Apply Rhizolizer™ at 1.5-3.0 fl. oz./acre.

Do not apply as foliar spray.

Product should be used within 2-3 hours of mixing the final application solution. Product can be co-applied with soil applied fertilizer solutions.

Conduct a jar test to ensure compatibility between products if Rhizolizer™ is tank mixed with fertilizer.

Do not tank mix with or apply directly after agricultural fungicides or anti-microbials.

Trees/Citrus: Apply 1.5 fl. oz. per grove acre through irrigation system. Fill irrigation injection system to at least 50% water volume, add appropriate amount of product, and fill tank with remainder of volume. Irrigate with water for 30 minutes after soil application.

Citrus Resets: Drench application is 0.01 fl. oz./tree or 5.5 fl. oz. per 35 gallons delivered at 8-10 fl. oz. per tree by metered sprayer. Add product to mixing tank once the tank is half full with water.

Vegetables/Fruit Crops: Apply product through drip or micro-sprinkler systems to soil at a rate of 3.0 fl. oz./acre in water. Apply 3.0 fl. oz./acre through irrigation system. Fill irrigation injection system to at least 50% water volume, add appropriate amount of product, and fill tank with remainder of volume. Irrigate with water for 30 minutes after soil application.

Corn: Mix with liquid fertilizer at standard rate for equivalent of 3-4 fl. oz./acre of Rhizolizer™.

Sod: Apply product through conventional boom spray or pivot at a rate of 3.0 fl. oz./acre. Fill spray tank to at least 50% water volume, add appropriate amount of product, and fill tank with remainder of volume.

For best activity from Rhizolizer™, apply in a tank-mixture with a kelp and a humic acid product. See those product labels for application details.

DIRECTIONS OR GENERAL RECOMMENDATIONS

Hazard Symbol: None **Signal Word:** None

Hazard Statement: This substance does not meet the criteria for classification as a dangerous substance.

Prevention Statements: Wear eye goggles and gloves. Avoid breathing spray.

Response: Wash hands after handling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed in storage or disposal.

Storage: Keep refrigerated and store out of reach of children. Store in original containers only. Keep out of direct sunlight.

Disposal: Dispose of all waste in accordance with local, state, and federal regulations. Triple rinse all used containers and puncture and dispose of in accordance with federal, state, and local law.

Spills: Clean up with spill kit, wash away excess fluids, and dispose of collected solution according to federal, state, and local laws.

CONDITION OF SALE & WARRANTY

Read the entire Conditions of Sale and Warranty before using this product. If these terms are unacceptable, return the unopened container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product reflect the opinions of experts based on field use and laboratory testing. Always follow label instructions. Injury to the crop on which the product is applied may result from the occurrence of unusual weather conditions, or the failure to follow the label directions or good application practices, all of which are /beyond the control of the seller. In addition, failure to follow label directions may cause injury to other crops, animals, humans, or the environment. All such risks are assumed by the buyer.

DISCLAIMER OF WARRANTIES:

NEITHER manufacturer Nor seller (the "providers") makes ANY express or implied warranty, AND BOTH providers DISCLAIM THE IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. IN NO CASE SHALL either provider be liable to the buyer or user of this product for any INCIDENTAL, consequential OR special damages. NO AGENT OR EMPLOYEE OF a Provider IS AUTHORIZED TO make any warranty or MODIFY THIS disclaimer.

LIMITATIONS OF LIABILITY:

THE EXCLUSIVE REMEDY OF BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID.