

Bombigro™

All Crops

FOLIAR TREATMENT

PRODUCT DESCRIPTION

Bombigro may be used as part of a foliar program to enhance photosynthesis, plant vigor and crop yield. Bombigro augments a balanced crop nutrition program. **For commercial agriculture use only.**

DIRECTIONS

Directions for use: Agitate the product to ensure it is uniform.
Application rate is one (1) gallon per acre. See table for crop specific application timing.

Application Rates

Crop	Application Timing
Alfalfa	After every cutting
Corn	Once - at V5
Cotton	Once - just prior to bloom
Peanut	Twice - at first flower and 14 days later
Soybean	Once - at R3
Wheat/Cereals	Once - after flag leaf has emerged

LOT #:

Exp Date: 01/01/2025



CONTAINS NONPLANT FOOD INGREDIENTS

1.0% Sophorolipid

1.0% Citric Acid

98% Total Other Ingredients (Inert)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

PRODUCT HANDLING

Storage: Keep out of direct sunlight. Do not exceed storage in 77°F and above for an extended period of time. Do not freeze product.

Disposal: Dispose of all waste in accordance with local, state, and federal regulations.

KEEP OUT OF REACH OF CHILDREN

No hazard symbol. This substance does not meet criteria for classification as a dangerous substance. Wear eye goggles and gloves.



Scan to see
Condition of
Sale &
Warranty

F002455

Emergency number:
Chemtrec (800) 424-9300

General Information:
Locus AG (888) 331-5008

Bombigro™ and the Locus AG logos are trademarks of Locus Agricultural Solutions

Net Content: 250 gal (946.3 L)
Density: 8.3 lb/gal (2.2 lb/L)

Manufactured by:
Locus Agricultural Solutions
30600 Aurora Rd., Ste 180, Solon, OH, 44139

Reformulation is prohibited. Copyright © 2024 All Rights Reserved